

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

6.436J/15.085J

Fall 2018

Problem Set 1

---

**Readings:**

- (a) Notes from Lecture 1.
- (b) Handout on background material on sets and real analysis (Recitation 1).

**Supplementary readings:**

- [C], Sections 1.1-1.4.
- [GS], Sections 1.1-1.3.
- [W], Sections 1.0-1.5, 1.9.

**Exercise 1.**

- (a) Let  $\mathbb{N}$  be the set of positive integers. A function  $f : \mathbb{N} \rightarrow \{0, 1\}$  is said to be *periodic* if there exists some  $N$  such that  $f(n + N) = f(n)$ , for all  $n \in \mathbb{N}$ . Show that the set of periodic functions is countable.
- (b) Does the result from part (a) remain valid if we consider rational-valued periodic functions  $f : \mathbb{N} \rightarrow \mathbb{Q}$ ?

**Exercise 2.** Let  $\{x_n\}$  and  $\{y_n\}$  be real sequences that converge to  $x$  and  $y$ , respectively. Provide a formal proof of the fact that  $x_n + y_n$  converges to  $x + y$ .

**Exercise 3.** We are given a function  $f : A \times B \rightarrow \mathbb{R}$ , where  $A$  and  $B$  are nonempty sets.

- (a) Assuming that the sets  $A$  and  $B$  are finite, show that

$$\max_{x \in A} \min_{y \in B} f(x, y) \leq \min_{y \in B} \max_{x \in A} f(x, y).$$

- (b) For general nonempty sets (not necessarily finite), show that

$$\sup_{x \in A} \inf_{y \in B} f(x, y) \leq \inf_{y \in B} \sup_{x \in A} f(x, y).$$

**Exercise 4.** A probabilistic experiment involves an infinite sequence of trials. For  $k = 1, 2, \dots$ , let  $A_k$  be the event that the  $k$ th trial was a success. Write down a set-theoretic expression that describes the following event:

$B$ : For every  $k$  there exists an  $\ell$  such that trials  $k\ell$  and  $k\ell^2$  were both successes.

*Note:* A “set theoretic expression” is an expression like  $\bigcup_{k>5} \bigcap_{\ell<k} A_{k+\ell}$ .

**Exercise 5.** Let  $f_n, f, g : [0, 1] \rightarrow [0, 1]$  and  $a, b, c, d \in [0, 1]$ . Derive the following set theoretic expressions:

(a) Show that

$$\{x \in [0, 1] \mid \sup_n f_n(x) \leq a\} = \bigcap_n \{x \in [0, 1] \mid f_n(x) \leq a\},$$

and use this to express  $\{x \in [0, 1] \mid \sup_n f_n(x) < a\}$  as a countable combination (countable unions, countable intersections and complements) of sets of the form  $\{x \in [0, 1] \mid f_n(x) \leq b\}$ .

(b) Express  $\{x \in [0, 1] \mid f(x) > g(x)\}$  as a countable combination of sets of the form  $\{x \in [0, 1] \mid f(x) > c\}$  and  $\{x \in [0, 1] \mid g(x) < d\}$ .

(c) Express  $\{x \in [0, 1] \mid \limsup_n f_n(x) \leq c\}$  as a countable combination of sets of the form  $\{x \in [0, 1] \mid f_n(x) \leq c\}$ .

(d) Express  $\{x \in [0, 1] \mid \lim_n f_n(x) \text{ exists}\}$  as a countable combination of sets of the form  $\{x \in [0, 1] \mid f_n(x) < c\}$ ,  $\{x \in [0, 1] \mid f_n(x) > c\}$ , etc. (Hint: think of  $\{x \in [0, 1] \mid \limsup_n f_n(x) > \liminf_n f_n(x)\}$ ).

**Exercise 6. Optional — not to be graded.**

This exercise develops an example that is meant to illustrate the following: if we work with fields instead of  $\sigma$ -fields, and if we only require finite additivity, then countable additivity will not be an automatic consequence, and the model may not correspond to any intuitive notion of probabilities.

Let  $\mathbb{N} = \mathbb{N}$  (the positive integers), and let  $\mathcal{F}_0$  be the collection of subsets of  $\mathbb{N}$  that either have finite cardinality or their complement has finite cardinality. For any  $A \in \mathcal{F}_0$ , let  $\mathbb{P}(A) = 0$  if  $A$  is finite, and  $\mathbb{P}(A) = 1$  if  $A^C$  is finite.

(a) Show that  $\mathcal{F}_0$  is a field but not a  $\sigma$ -field.

(b) Show that  $\mathbb{P}$  is finitely additive on  $\mathcal{F}_0$ ; that is, if  $A, B \in \mathcal{F}_0$ , and  $A, B$  are disjoint, then  $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$ .

- (c) Show that  $\mathbb{P}$  is not countably additive on  $\mathcal{F}_0$ ; that is, construct a sequence of disjoint sets  $A_i \in \mathcal{F}_0$  such that  $\cup_{i=1}^{\infty} A_i \in \mathcal{F}_0$  and  $\mathbb{P}(\cup_{i=1}^{\infty} A_i) \neq \sum_{i=1}^{\infty} \mathbb{P}(A_i)$ .
- (d) Construct a decreasing sequence of sets  $A_i \in \mathcal{F}_0$  such that  $\cap_{i=1}^{\infty} A_i = \emptyset$  for which  $\lim_{i \rightarrow \infty} \mathbb{P}(A_i) \neq 0$ .

MIT OpenCourseWare  
<https://ocw.mit.edu>

6.436J / 15.085J Fundamentals of Probability  
Fall 2018

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>