

Markov processes – II

- review and some warm-up
 - definitions, Markov property
 - calculating the probabilities of trajectories
- steady-state behavior
 - recurrent states, transient states, recurrent classes
 - periodic states
 - convergence theorem
 - balance equations
- birth-death processes


review

- discrete time, discrete state space, time-homogeneous
 - transition probabilities
 - Markov property


- $$\begin{aligned} r_{ij}(n) &= \mathbf{P}(X_n = j \mid X_0 = i) \\ &= \mathbf{P}(X_{n+s} = j \mid X_s = i) \end{aligned}$$

- key recursion:

$$r_{ij}(n) = \sum_{k=1}^m r_{ik}(n-1)p_{kj}$$



warmup




$$\mathbf{P}(X_1 = 2, X_2 = 6, X_3 = 7 \mid X_0 = 1) =$$

$$\mathbf{P}(X_4 = 7 \mid X_0 = 2) =$$

review: recurrent and transient states




- state i is **recurrent** if “starting from i , and from wherever you can go, there is a way of returning to i ”
- if not recurrent, called **transient**




- **recurrent class**: a collection of recurrent states communicating only between each other

periodic states in a recurrent class


The states in a recurrent class are periodic if they can be grouped into $d > 1$ groups so that all transitions from one group lead to the next group



steady-state probabilities




- does $r_{ij}(n) = \mathbf{P}(X_n = j \mid X_0 = i)$ converge to some π_j ?
- theorem: yes, if:
 - recurrent states are all in a single class, and
 - single recurrent class is not periodic



- assuming “yes”, start from key recursion $r_{ij}(n) = \sum_k r_{ik}(n-1)p_{kj}$
 - take the limit as $n \rightarrow \infty$
- $$\pi_j = \sum_k \pi_k p_{kj}$$
- need also: $\sum_j \pi_j = 1$

example




$$\pi_j = \sum_k \pi_k p_{kj}$$

visit frequency interpretation


- balance equations

$$\pi_j = \sum_k \pi_k p_{kj}$$



- (long run) frequency of being in j : π_j






- frequency of transitions $1 \rightarrow j$: $\pi_1 p_{1j}$



- frequency of transitions into j : $\sum_k \pi_k p_{kj}$





birth-death processes I



$$\pi_i p_i = \pi_{i+1} q_{i+1}$$

birth-death processes II



$$\pi_i p_i = \pi_{i+1} q_{i+1}$$

$$\sum_j \pi_j = 1$$

special case: $p_i = p$ and $q_i = q$ for all i

$$\rho = p/q \quad \pi_{i+1} = \pi_i \frac{p}{q} = \pi_i \rho$$

$$\pi_i = \pi_0 \rho^i \quad i = 0, 1, \dots, m$$

- assume $p = q$

- assume $p < q$ and $m \approx \infty$

$$\pi_0 = 1 - \rho \quad \mathbf{E}[X_n] = \frac{\rho}{1 - \rho} \text{ (in steady-state)}$$

MIT OpenCourseWare
<https://ocw.mit.edu>

Resource: Introduction to Probability
John Tsitsiklis and Patrick Jaillet

The following may not correspond to a particular course on MIT OpenCourseWare, but has been provided by the author as an individual learning resource.

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.